



Encouraging Physical Activity in Diabetes Patients Through Automatic Personalized Feedback Via Reinforcement Learning Improves Glycemic Control

Irit Hochberg,¹ Guy Feraru,²
Mark Kozdoba,³ Shie Manor,³
Moshe Tennenholtz,⁴ and
Elad Yom-Tov⁵

Diabetes Care 2016;39:e1–e2 | DOI: 10.2337/dc15-2340

Most patients with type 2 diabetes are sedentary despite the clear benefit of regular physical activity, including better glucose control and improvement in quality of life (1). Smartphones could potentially improve patient care by continual communication with patients and sensors that quantify patient behavior (2). Nevertheless, the use of personalized messages that take into account the actual behavior of patients and learn to reinforce it has not yet been reported.

We provided a total of 27 patients with type 2 diabetes who did not perform regular physical activity prior to recruitment with a pedometer installed on their personal smartphones and a personal plan for physical activity. Patients were randomized into a control group ($n = 7$) and a treatment group ($n = 20$), and received textual messages (via the Short Message Service on the smartphone) one to seven times a week to encourage physical activity. In the treatment group, the messages comprised positive feedback (with and without a social component) and negative feedback. Messages to patients in the treatment group were initially selected through nonchanging expert-generated rules (“initial policy”) and later were personalized through an automatic Reinforcement

Learning (3) algorithm (“personalized policy”), which learns to optimize messages to improve each participant’s compliance with the activity regimen. Patients in the control group received constant weekly reminders to exercise (“control”). Follow-up HbA_{1c} tests were performed every 3 months. Physicians were blinded to the randomization.

Participants who received messages tailored by the personalized policy increased the amount of activity (e.g., walking) and pace of walking over time (as seen in the positive slopes of the graphs of these variables), while the control group patients did not (Table 1). Allocation to the personalized policy, higher initial HbA_{1c} level, and lower activity targets led to a superior reduction in HbA_{1c} levels ($R^2 = 0.405$, $P < 0.0001$). In a questionnaire, patients in the treatment group reported that the messages helped them to increase ($P = 0.01$) and to maintain ($P = 0.07$) physical activity, while control patients reported that messages were ineffective.

The learning algorithm improved gradually in predicting which messages would lead participants to exercise. On average, the best daily message was a positive-feedback message with a social component (average improvement

of 8.8% in activity in the day following such a message), and the best consecutive messages were a positive social message after a negative-feedback message (42.7% improvement). The least effective message was a positive-feedback message without social reference (9.9% reduction), and the least effective consecutive messages were a negative-feedback message after a positive social message (–61.4%). We also clustered participants by their response to the different types of messages and found that patients can be divided into the following three groups: one that reacted negatively to any message, one that only reacted positively to the positive—with social component message, and a third where patients reacted positively to all messages, especially a positive social message or positive self-message. This demonstrates the importance of individually tailored feedback as delivered by our algorithm.

These results suggest that a mobile phone application with a learning algorithm can improve adherence to exercise in patients with diabetes. Because a personalized learning algorithm is automated, it can be used in large populations to improve health and glycemic control.

Q:1

Q:2

e-LETTERS – OBSERVATIONS

¹Rambam Health Care Campus, Haifa, Israel

²Faculty of Medicine, Technion—Israel Institute of Technology, Haifa, Israel

³Faculty of Electrical Engineering, Technion—Israel Institute of Technology, Haifa, Israel

⁴Faculty of Industrial Engineering and Management, Technion—Israel Institute of Technology, Haifa, Israel

⁵Microsoft Research, Herzliya, Israel

Corresponding Author: Irit Hochberg, i_hochberg@rambam.health.gov.il.

Clinical trial reg. no. NCT02612402, clinicaltrials.gov.

© 2016 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered.

Q:3

Table 1—Rates of improvement in physical activity performed and in the rate of walking

	Control (N = 7)	Treatment (N = 20)		P value (control vs. personalized) ^a
		Initial	Personalized	
Demographics				
Female sex	1	8		0.36
Age, years ^b	55.1 ± 3.6	58.7 ± 2.1		0.56
Initial HbA _{1c} % (mmol/mol)	8.7 (72)	7.7 (61)		1.00
Outcomes^c				
Slope of the change in activity, min of walking/day over time	−0.004 (0.002)	−0.001 (0.008)	0.012 (0.002)	2 * 10 ^{−5}
Slope of the rate of walking, Hz/day over time	−0.010 (0.007)	−0.009 (0.005)	0.002 (0.005)	0.04

Initial policy refers to a rule-based policy for sending messages. Personalized policy refers to messages that were optimized using the learning algorithm to maximize individual activity. The slope of change in activity is measured by a linear fit to the plotted amount of daily exercise over time. The slope of the rate of walking is the change in the number of steps per minute during walking over time. ^aThe P value was calculated by t test.

^bValues are reported as the mean ± SEM. ^cValues in parentheses are the SEM.

Acknowledgments. The authors thank all the participants for generously helping us in this study.

Duality of Interest. All authors have completed the Unified Competing Interest form. No potential conflicts of interest relevant to this article were reported; all authors preformed this study as part of their salaried employment within their respective institutions. The authors have no financial relationships with any organizations that might have an interest in the submitted work in the previous 3 years, and no other relationships or activities that could appear to have influenced the submitted work.

Author Contributions. I.H. conceived the idea, designed the study, obtained institutional review board approval, recruited the patients, collected data, and wrote the article. G.F. collected and analyzed the medical data. M.K. designed and built the app. S.M. helped to design the study and the algorithm, and designed and built the app. M.T. helped to design the study and the algorithm. E.Y.-T. developed the idea, designed and implemented the study, analyzed the results, and wrote the article. I.H. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

References

1. Pronk NP, Remington PL; Community Preventive Services Task Force. Combined diet and physical activity promotion programs for prevention of diabetes: Community Preventive Services Task Force recommendation statement. *Ann Intern Med* 2015;163:465–468
2. de Jongh T, Gurol-Urganci I, Vodopivec-Jamsek V, Car J, Atun R. Mobile phone messaging for facilitating self-management of long-term illnesses. *Cochrane Database Syst Rev* 2012;12:CD007459
3. Gatti C. Reinforcement learning. In *Design of Experiments for Reinforcement Learning*. New York, Springer, 2014, p. 7–52